




Transient phenomena


Miguel Pérez-Torres (IAA-CSIC, Granada)

(includes also contributions from
G. Ghirlanda, M. Giroletti, J. Miller-Jones,
T. O'Brien, Z. Paragi)





The transient parameter space



- 
- Pulsars
 - FRBs
 - CCSNe/SNe Ia
 - GRBs
 - XRBs/quasars
 - ULX
 - Novae/CVs
 - AGNs, TDEs
 - GW follow-up


FRBs

- A tool to study the cosmic web: the distribution of matter in the Universe
- Most of this matter within the galaxies is invisible otherwise




Fast Radio Bursts (FRBs)


FRB localization is key



5-ms image (dispersion corrected) of one burst.






Variable steady source associated to FRB 150418 (Giroletti+16)



Localisation of the repeating FRB 121102 (Marcote+17)

GW170817 EM counterpart


Abbott+17



Structured jet and cocoon models tested via light curve modelling

VLBI obs-ns can measure expansion of ejecta


CCSNe



Expansion of SN2008iz in M82 imaged with VLBI (Brunthaler+2010)

Type Ia SNe

- What are their progenitors?
 - Single degenerate (SD) channel => Prompt radio emission
 - Double-Degenerate (DD) channel => No prompt radio emission



(Pérez-Torres+2015)

Type Ia SNe

5.0 GHz Continuum MERLIN Observations of the Type Ia SN 2013dy

ATel #5619; *M. Perez-Torres (IAA-CSIC/CEFCA, Spain), M. Argo (JBCA, Manchester), P.*

Lu
Bjo **EVN measurements show no evidence for radio emission from the Type Ia SN 2014J**

JBCA), C. I. xford/Soton),

ATel #6153; *M. Perez-Torres (IAA-CSIC, Granada; CEFCA, Teruel), P. Lundqvist (Dept. of*

Subj
Astrono
Astronomy
Alber
Manches
Mart&iacu
(Max-Planck
Tight constraints on the mass-loss rate of the Type Ia SN 2016coj with e-MERLIN

ATel #10168; *M. Perez-Torres (IAA-CSIC), P. Lundqvist, E. Kundu (Stockholm), J. Moldon*
(JBCA), C. I. Bjornsson (Stockholm University), C. Fransson (Stockholm University)

Radio constraints on the mass-loss rate of the Type Ia SN 2018gv

We
10.4
CBE
reac
obse
Defi
MH:

Subjects: Radio
Referred to by ATel

ATel #11211; *S. D. Ryder (AAO), P. Lundqvist (Stockholm University), M. A. Perez-Torres (IAA-CSIC), E. Kundu (Stockholm University), C. I. Bjornsson (Stockholm University)*

Mass-loss rate constraints on the Type Ia SN 2018pv from e-MERLIN observations

our
DEC
this
We
regio
limit
uppe
(3-si

We report d
2014J, whic
<http://www.cbet.org>
CBET #379
a frequency
Effelsberg,
Noto and Si
bandwidth;
synthesized
observations
et al. (ATel
standard rad

We report e-MERLIN on 28.18 May 2018 (MJD #9095). Our observations were carried out with the e-MERLIN radio telescopes (Cambridge, Pickering, and Jodrell Bank) at a frequency of 1.51 GHz and a beam of (0.13 x 0.13) arcsec. This is the first radio detection of SN 2018gv (J2000.0 05h 16m 16.1s, +36° 59' 11.6").

Subjects: Radio, Supernovae, Transient

Referred to by ATel #: 11324

[Tweet](#) [Recommend 4](#)

as in
We
Type

We find no radio emission at 5.0 GHz, in the region surrounding the supernova. This corresponds to an upper limit of 1.67 erg/s/Hz and 1.67 microJy/beam at 1.67 GHz. Assuming a progenitor mass of 1.9 solar masses per year (3-sigma), this model, our data are consistent with the progenitor of 1.9 solar masses per year (3-sigma).

The young Type Ia SN 2018gv (ATel #11324) was discovered on 20 Feb 2018 (MJD 58153.13) in the nearby (z=0.0031) galaxy NGC3941 (Tsuboi, TNS discovery report #16800), and a spectrum on 8.78 February 2018 (MJD 58158.78) confirmed the SN as a Type Ia event a few days before maximum (Yamanaka, TNS classification report #1712). Our observations were carried out on 9-10 February 2018 UT (MJD 58159.08), six days after the SN discovery. We centered our observations at the position of the optical discovery (J2000.0 05h 16m 16.1s, +36° 59' 11.6"). We find


Subjects: Radio, Supernovae

[Tweet](#) [Recommend 1](#)

We observed the young Type Ia supernova 2018pv with the electronic Multi-Element Radio Linked Interferometer Network (e-MERLIN) at 5.1 GHz. SN 2018pv was discovered on 3.63 February 2018 UT (MJD 58153.13) in the nearby (z=0.0031) galaxy NGC3941 (Tsuboi, TNS discovery report #16800), and a spectrum on 8.78 February 2018 (MJD 58158.78) confirmed the SN as a Type Ia event a few days before maximum (Yamanaka, TNS classification report #1712). Our observations were carried out on 9-10 February 2018 UT (MJD 58159.08), six days after the SN discovery. We centered our observations at the position of the optical discovery (J2000.0 05h 16m 16.1s, +36° 59' 11.6"). We find

on 20 Feb 2018; 14:34 UT
Credential Certification: Miguel A. Perez-Torres (torres@iaa.es)

Type Ia SN progenitors - SD channel



**SKA chapter on SNe (Pérez-Torres+2015).
Plot adapted from Pérez-Torres+2014**

GRBs




Figure from Nakar & Piran (1993)

VLBI obs-ns extremely useful

=> **Deep flux measurements + resolution**

- Jet properties (structure, dynamics, orientation)
- Shock properties (e.g. energy spectrum of e-),
- Environment (ISM, wind)

GRBs



GRB 030329: the best ever radio characterization (bright and close)

VLBI observations crucial to disentangle GRB environment

Global VLBI obs-ns ($t=5.5$ yr!)



- Single power-law decay ($t^{-1.27}$)
- Proper motion < 0.067 mas yr $^{-1}$
- Size evolution

- Jet seen close to the LOS
- Expansion in the ISM
- Emission due to external shock, accelerated electrons ($p=2.5$)





GRBs

GRB151027A




GRBs - GRB 171205A




- VLBA, EVN, and eMERLIN obs-ns
- Discern Cannonball/Fireball model
- Discern ISM/wind interaction

X-ray binaries



XTE J1908+094 expanding jet (Rushton+17)



Cyg X-3 giant flare (Egron+17)

Novae

White dwarf + Main Sequence/Red Giant companion



- Outbursts due to thermonuclear runaway in accreted material on WD surface.
- The white dwarf is not destroyed and another nova outburst may occur ~ 1 to 1000's of years later.

Novae


Gamma-ray nova V959 Mon (Chomiuk et al 2014)

Expanding non-thermal components
seen with EVN

- High-resolution radio imaging of their expanding aspherical remnants to understand their ejection geometry, including jets
- Combining radio, X-ray and gamma-ray observations to understand role of shocks in particle acceleration
- Understanding explosions on massive WDs and their link to Type Ia SNe



e-MERLIN imaging of V959 Mon (Healy et al 2017)




Real-time jet evolution

- **How relativistic are XRB jets?**
 - Accurate distances essential
- **How do jets couple to accretion flow?**
 - VLBI proper motions give ejection time
 - Tie to X-ray spectral / timing signatures

18 months


Gomez et al. (2001)



4 hours
Miller-Jones et al. (2018)

Fender et al. (1999)

11 days










Precision astrometry

Fundamental physical parameters

- XRB jet cores provide astrometric targets
 - *Proper motion*: Natal kicks and BH formation
 - *Parallax*: Model-independent distances
 - *Residuals*: Jet size scales
 - *Orbital motion*: Component masses






BH natal kick distribution needed to simulate LIGO merger progenitors

Nuclear Transients

An extremely prolific SN factory in Arp 299-A revealed with the eEVN


- Rich cluster of compact radio sources in the nuclear region of Arp299A
- SNe and/or SNRs, likely embedded in SSCs.
- Evidence of recent RSNs
- Radio emission levels typical of Type II SNe

(Pérez-Torres+2009, A&A Letters)



The nuclear starburst in Arp 299A


- >26 sources detected
- CCSNe and SNRs
- AGN unveiled
- Evidence for new SNe
- CCSN ≥ 0.8 SN/yr
- Flickering microQSO



10 pc

Pérez-Torres+09
Pérez-Torres+10
Bondi & PérezTorres+12)

Tidal Disruption Events (TDEs)





ASSASN-14li resolved at pc-scales with the EVN (Romero-Cañizales+2016)

Source nature unclear:


- Core-relativistic jet?
- Core-non-relativistic jet?
- BBH?

Tidal Disruption Events (TDEs)



No apparent superluminal motion in Sw J1644+5734 unveiled with the EVN (Yang+2016)

Discovery of an extremely luminous nuclear outburst in Arp 299B I




- Systematic near-IR search for nuclear SNe in starburst galaxies using the WHT
- Discovery of an extremely luminous nuclear outburst in the near-IR in Jan. 2005
- **Only detected in the IR, in optical completely obscured** by interstellar dust
- Near-IR (JHKs) follow-up from the WHT, NOT, Gemini-N, mid-IR from Spitzer




Radio evolution from VLBI observations

- New compact source detected in Jul 2005
- Developed prominent extended, jet-like structure by 2011
- EVN observations crucial




A TDE with a resolved, relativistic radio jet in a galaxy merger



- Radio observations resolved expanding, decelerating jet
- Probe jet formation around a SMBH
- Dust reprocessing may explain differences betw theo and observed TDE lumins


AT 2017gbl in IRAS 23436+5257




- Near-IR nuclear burst detected in July 2017
- IR properties similar to the Arp299B-AT1 event

Kool et al. (2017)

VLBI obs-ns of AT 2017gbl in the nuclear region of IRAS 23436+5257



Peak = 880 uJy/beam



Peak = 1720 uJy/beam

- $L_{\text{radio}} \sim 3.2e38 \text{ erg/s}$
- Inverted spectral index ($\alpha = 1.1$; $S_{\nu} \sim \nu^{\alpha}$)
- Compatible with a LLAGN. It could also be AT 2017gbl

Pérez-Torres et al. (2017, ATel)

Prospects for transients

- **“Prompt” emission**
 - Probe self-absorbed events in early phases of GRBs, GW events, TDEs, AGN flares (=> higher freq obs-ns)
 - Get early measurements as references, both for structure (motions/expansions of novae, SNe, jets,...) and light curve (initial flux density, or upper limit)
- **“Late” emission**
 - Follow evolution on different spatial scales and with full spectral information
 - iteratively refine models and design observations to pick sources at most suitable times
- **Exploit sensitivity** through large apertures and wide bandwidth.

(Some) issues

- **Alert/triggering procedures** - not straightforward to trigger fast repointing on different sites, particularly if not dedicated/full time arrays
 - Different transients require different triggers (from space or ground, photometry or spectrum) and different reaction times
- Disk availability and shipping, correlation time can cause delays - **real-time VLBI** is still a **relatively scarce resource**
- **Arrays** - Small, flexible arrays for prompt observations; full/global arrays for follow up of truly interesting events?
- **Calibration uncertainties** should go down; overall performance and reliability of the arrays
- **SKA I (and precursors) need to be fully VLBI-ready** (receivers/backends compatible with existing telescopes, phased array capabilities, data formats and transfer, ...)