




# Transient phenomena

Miguel Pérez-Torres (IAA-CSIC, Granada)


(includes also contributions from  
G. Ghirlanda, M. Giroletti, J. Miller-Jones,  
T. O'Brien, Z. Paragi)




EVN Vision f2f meeting, March 1st, 2018, Zaandam



# The transient parameter space





Updated plot of Cordes+2004

- 
- Pulsars
  - FRBs
  - CCSNe/SNe Ia
  - GRBs
  - XRBs/quasars
  - ULX
  - Novae/CVs
  - AGNs, TDEs
  - GW follow-up


# FRBs

- A tool to study the cosmic web: the distribution of matter in the Universe
- Most of this matter within the galaxies is invisible otherwise




# Fast Radio Bursts (FRBs)


FRB localization is key



5-ms image (dispersion corrected) of one burst.






Variable steady source associated to FRB 150418 (Giroletti+16)



Localisation of the repeating FRB 121102 (Marcote+17)

# GW170817 EM counterpart


Abbott+17



Structured jet and cocoon models tested via light curve modelling

VLBI obs-ns can measure expansion of ejecta


# CCSNe



Expansion of SN2008iz in M82 imaged with VLBI (Brunthaler+2010)

# Type Ia SNe

- What are their progenitors?
  - Single degenerate (SD) channel => Prompt radio emission
  - Double-Degenerate (DD) channel => No prompt radio emission



(Pérez-Torres+2015)



# Type Ia SNe

## 5.0 GHz Continuum MERLIN Observations of the Type Ia SN 2013dy

ATel #5619; *M. Perez-Torres (IAA-CSIC/CEFCA, Spain), M. Argo (JBCA, Manchester), P.*

*Lu Bjo* **EVN measurements show no evidence for radio emission from the Type Ia SN 2014J**

*JBCA), C. I. xford/Soton),*

ATel #6153; *M. Perez-Torres (IAA-CSIC, Granada; CEFCA, Teruel), P. Lundqvist (Dept. of*

*Subj Astronomy Alber Manches Mart&iacu (Max-Planck)* **Tight constraints on the mass-loss rate of the Type Ia SN 2016coj with e-MERLIN**

ATel #10168; *M. Perez-Torres (IAA-CSIC), P. Lundqvist, E. Kundu (Stockholm), J. Moldon (JBCA, Manchester), S. Ryder (Maccquarie University/AAO)*

**Radio constraints on the mass-loss rate of the Type Ia SN 2018gv**

ATel #11211; *S. D. Ryder (AAO), P. Lundqvist (Stockholm University), M. A. Perez-Torres (IAA-CSIC), E. Kundu (Stockholm University), C. Fransson (Stockholm University)*  
*on Credential Certif*

**Mass-loss rate constraints on the Type Ia SN 2018pv from e-MERLIN observations**

ATel #11324; *M. Perez-Torres (IAA-CSIC, Granada), P. Lundqvist (Stockholm University), J. Moldon (JBCA, Manchester), S. Ryder (Maccquarie University/AAO), E. Kundu (Stockholm University), E. Varenus (JBCA, Manchester), A. Alberdi (IAA-CSIC, Granada), R. Beswick (JBCA, Manchester), C.-I. Bjornsson (Stockholm University), C. Fransson (Stockholm University)*

*on 20 Feb 2018; 14:34 UT*

*Credential Certification: Miguel A. Perez-Torres (torres@iaa.es)*

Subjects: Radio, Supernovae, Transient

Referred to by ATel #: 11324

Tweet Recommend 4

Subjects: Radio, Supernovae

Subjects: Radio, Supernovae

Tweet Recommend 1

We observed the young Type Ia supernova 2018pv with the electronic Multi-Element Radio Linked Interferometer Network (e-MERLIN) at 5.1 GHz. SN 2018pv was discovered on 3.63 February 2018 UT (MJD 58153.13) in the nearby ( $z=0.0031$ ) galaxy NGC3941 (Tsuboi, TNS discovery report #16800), and a spectrum on 8.78 February 2018 (MJD 58158.78) confirmed the SN as a Type Ia event a few days before maximum (Yamanaka, TNS classification report #1712). Our observations were carried out on 9-10 February 2018 UT (MJD 58159.08), six days after the SN discovery. We centered our observations at the position of the optical discovery (J2000.0 coordinates RA=11:52:55.70, DEC=+36:59:11.60, TNS discovery report #16800). We find

We report d  
our 2014J, whic  
DEC http://www.  
this CBET #379  
We find no e  
GHz, in the  
erg/s/Hz and  
at 1.67 GHz  
limits to the  
for the prog  
published in

Subjects: Ra  
Referred to

Referred to by ATel  
 Tweet Reco

We report d  
our 2014J, whic  
DEC http://www.  
this CBET #379

We find no e  
GHz, in the  
erg/s/Hz and  
at 1.67 GHz  
limits to the  
for the prog  
published in

Subjects: Ra  
Referred to

Referred to by ATel  
 Tweet Reco

We report d  
our 2014J, whic  
DEC http://www.  
this CBET #379

We find no e  
GHz, in the  
erg/s/Hz and  
at 1.67 GHz  
limits to the  
for the prog  
published in

Subjects: Ra  
Referred to

Referred to by ATel  
 Tweet Reco

Subjects: Supernovae

Referred to by ATel

Tweet Reco

We report e-MERL  
on 28.18 May 201  
#9095). Our observ  
hr, one week after t

radio telescopes  
(Cambridge, Pickm  
of 1.51 GHz and u  
beam of (0.13 x 0.  
discovery ( J2000.

(16x16) sq. arcsec  
the region surroun  
corresponds to an  
sigma). Assuming t  
Torres et al. (2014,  
very stringent uppe  
masses per year (3-

this model, our dat  
MERLIN staff for  
supernovae, aimed  
research leading to  
Framework Programme (FP/2007-2


Subjects: Radio, Supernovae, Transient

Referred to by ATel #: 11324

Tweet Recommend 4

Subjects: Radio, Supernovae

# Type Ia SN progenitors - SD channel



**SKA chapter on SNe (Pérez-Torres+2015).  
Plot adapted from Pérez-Torres+2014**

# GRBs




Figure from Nakar & Piran (1993)

## VLBI obs-ns extremely useful

=> **Deep flux measurements + resolution**

- Jet properties (structure, dynamics, orientation)
- Shock properties (e.g. energy spectrum of e-),
- Environment (ISM, wind)

# GRBs



GRB 030329: the best ever radio characterization (bright and close)

VLBI observations crucial to disentangle GRB environment

Global VLBI obs-ns ( $t=5.5$  yr!)



- Single power-law decay ( $t^{-1.27}$ )
- Proper motion  $< 0.067$  mas yr $^{-1}$
- Size evolution

- Jet seen close to the LOS
- Expansion in the ISM
- Emission due to external shock, accelerated electrons ( $p=2.5$ )





# GRBs

## GRB151027A




# GRBs - GRB 171205A




- VLBA, EVN, and eMERLIN obs-ns
- Discern Cannonball/Fireball model
- Discern ISM/wind interaction

# X-ray binaries



XTE J1908+094 expanding jet (Rushton+17)



Cyg X-3 giant flare (Egron+17)

# Novae

White dwarf + Main Sequence/Red Giant companion

- Outbursts due to thermonuclear runaway in accreted material on WD surface.
- The white dwarf is not destroyed and another nova outburst may occur  $\sim 1$  to 1000's of years later.





# Novae


Gamma-ray nova V959 Mon (Chomiuk et al 2014)

Expanding non-thermal components  
seen with EVN

- High-resolution radio imaging of their expanding aspherical remnants to understand their ejection geometry, including jets
- Combining radio, X-ray and gamma-ray observations to understand role of shocks in particle acceleration
- Understanding explosions on massive WDs and their link to Type Ia SNe



e-MERLIN imaging of V959 Mon (Healy et al 2017)




## Real-time jet evolution

- **How relativistic are XRB jets?**
  - Accurate distances essential
- **How do jets couple to accretion flow?**
  - VLBI proper motions give ejection time
  - Tie to X-ray spectral / timing signatures

18 months


Gomez et al. (2001)



4 hours  
Miller-Jones et al. (2018)

Fender et al. (1999)

11 days










# Precision astrometry

## Fundamental physical parameters

- XRB jet cores provide astrometric targets
  - *Proper motion*: Natal kicks and BH formation
  - *Parallax*: Model-independent distances
  - *Residuals*: Jet size scales
  - *Orbital motion*: Component masses






**BH natal kick distribution needed to simulate LIGO merger progenitors**

# Nuclear Transients

An extremely prolific SN factory in Arp 299-A revealed with the eEVN

- Rich cluster of compact radio sources in the nuclear region of Arp299A
- SNe and/or SNRs, likely embedded in SSCs.
- Evidence of recent RSNs
- Radio emission levels typical of Type II SNe

(Pérez-Torres+2009, A&A Letters)




# The nuclear starburst in Arp 299A

- >26 sources detected
- CCSNe and SNRs
- AGN unveiled
- Evidence for new SNe
- CCSN  $\geq 0.8$  SN/yr
- Flickering microQSO

10 pc

Pérez-Torres+09  
Pérez-Torres+10  
Bondi & PérezTorres+12)

# Tidal Disruption Events (TDEs)





ASSASN-14li resolved at pc-scales with the EVN (Romero-Cañizales+2016)

Source nature unclear:


- Core-relativistic jet?
- Core-non-relativistic jet?
- BBH?

# Tidal Disruption Events (TDEs)



No apparent superluminal motion in Sw J1644+5734 unveiled with the EVN (Yang+2016)

# Discovery of an extremely luminous nuclear outburst in Arp 299B I




- Systematic near-IR search for nuclear SNe in starburst galaxies using the WHT
- Discovery of an extremely luminous nuclear outburst in the near-IR in Jan. 2005
- **Only detected in the IR, in optical completely obscured** by interstellar dust
- Near-IR (JHKs) follow-up from the WHT, NOT, Gemini-N, mid-IR from Spitzer






# AT 2017gbl in IRAS 23436+5257




- Near-IR nuclear burst detected in July 2017
- IR properties similar to the Arp299B-AT1 event

Kool et al. (2017)

# VLBI obs-ns of AT 2017gbl in the nuclear region of IRAS 23436+5257



Peak = 880 uJy/beam



Peak = 1720 uJy/beam

- $L_{\text{radio}} \sim 3.2e38$  erg/s
- Inverted spectral index ( $\alpha = 1.1$ ;  $S_{\nu} \sim \nu^{\alpha}$ )
- Compatible with a LLAGN. It could also be AT 2017gbl

Pérez-Torres et al. (2017, ATel)

# VLBI prospects for transients

- **Target localization:** mas-precision and accuracy is a must
- **Target imaging:** Images at mas-scales much needed
- **Ultra-high sensitivity:** Needed both for imaging and localization purposes, as well as for detection of faint, diffuse emission. Currently a few microJy/b. Should aim at  $\sim 1$  microJy/b sensitivity, i.e., equal to SKA1-MID
- **Astrometric capabilities** of VLBI currently experience a renaissance, with strong implications in many fields

# VLBI prospects/issues for transients

- **Spectral index information:** Frequency agility. EVN still lacks it, whether as EVN-regular, or as eEVN
- **Target follow-up => Multiple frequencies, multiple visits;** VLBA often better suited than EVN; eEVN mitigates this issue
- **Need to image simultaneously at different angular scales:** => EVN+eMERLIN, VLA=eVLA(=VLA+VLBA),...
- **Calibration uncertainties must go down:** overall performance and reliability of arrays must get better, esp. the EVN.

# VLBI prospects/issues for transients

- **Alert/triggering procedures** - not straightforward to trigger fast **repointing** on different sites, particularly if not dedicated/full time arrays
  - Different transients require different triggers (from space or ground, photometry or spectrum) and different reaction times
- Disk availability and shipping, correlation time can cause delays - **real-time VLBI** is still a **relatively scarce resource**
- **Arrays** - Small, flexible arrays for prompt observations; full/global arrays for follow up of truly interesting events?

# Challenge for the EVN

Towards a 1 microJy/b sensitivity,  
frequency-agile,  
flexible, multi-scale VLBI array  
for the next decade